arXiv:1106.1795v2 [hep-th] 26 Mar 2012

4+1 dimensional homogeneous anisotropic string cosmological models

B. Mojaveri[*] and A. Rezaei-Aghdam[†]
Department of Physics, Azarbaijan university of Tarbiat Moallem, 53714-161, Tabriz, Iran .

January 28, 2013

Abstract

We present exact solutions of string cosmological models characterized by five dimensional metrics (with four dimensional real Lie groups as isometry groups), space independent dilaton and vanishing torsion. As an example we consider $VII_0 \oplus R$ model and show that it is equivalent to the $4+1$ dimensional cosmological model coupled to perfect fluid with negative deceleration parameters (accelerating universe).

---

[*] e-mail: bmojaveri@azaruniv.edu
[†] e-mail: rezaei-a@azaruniv.edu

1 Introduction

Some of the important problems of primordial cosmology such as, why is our universe four-dimensional? why does a small vacuum energy-density seem to survive until today after inflation? and other problems should find a satisfactory solution in the context of string cosmology [1]. The dynamics of the very early universe below the string scale may have been determined by the string effective action [2]. The idea that string theory, with its fundamental length scale, could resolve the big bang singularity by effective limiting space-time curvature has led, during the last decade, to a variety of models [3] where the big bang represent a turning (rather than an end)-point in the history of the early Universe (see also [4, 5]). The general, spatially isotropic and homogeneous, FRW string cosmology is known and well understood [6]. Furthermore, it is generally accepted that spatial anisotropy might have been important in the very early universe and the study of the string cosmology models that relax the FRW assumptions is therefore well motivated. The spatially homogeneous anisotropic Bianchi cosmologies [7, 8, 9] have been studied previously and some inhomogeneous generalizations of these models were presented [10]. Bianchi cosmologies admit a three-dimensional real Lie group of isometries that acts simply-transitively on three-dimensional, space like orbits [11]. Some generalization of these models to models with four-dimensional real Lie groups was also performed [12].

Here we present exact solutions of string cosmological models characterized by five dimensional metric (with four dimensional real Lie group as isometry groups), a dilaton field and vanishing torsion.

The paper is organized as follows. In section two, we review the homogenous anisotropic string cosmology in d-dimension [9] and then obtain main equations in $4+1$ dimension. After then in section three, we obtain metrics over real four dimensional Lie groups. Subsequently, in section four we solve and obtain exact solutions of equations of motions for $4+1$ dimensional homogenous anisotropic string cosmological models related to real four dimensional Lie algebras. Finally in section five we consider example $VII_0 \oplus R$ and show that it is equivalent with $4+1$ dimensional cosmological coupled to perfect fluid with negative decelerating parameters (accelerating universe). Some concluding remarks including T- duality discussions are given in section six.

2 Review of homogeneous anisotropic string cosmology

We start with string effective action [2]:

$$S_{eff} = \int d^{d+1}x\sqrt{-g}e^{\phi}(R - \frac{1}{12}H_{\mu\nu\rho}H^{\mu\nu\rho} + \partial_\mu\phi\partial^\mu\phi - \Lambda). \tag{1}$$

The equations of motion for this action have the following forms

$$R_{\mu\nu} - \frac{1}{4}H^2_{\mu\nu} - \nabla_\mu\nabla_\nu\phi = 0, \tag{2}$$

$$\nabla^2(e^{\phi}H_{\mu\nu\lambda}) = 0, \tag{3}$$

$$-R + \frac{1}{12}H^2 + 2\nabla^2\phi + (\partial_\mu\phi)^2 + \Lambda = 0, \tag{4}$$

where the contractions $H^2_{\mu\nu} = H_{\mu\kappa\lambda}H_\nu{}^{\kappa\lambda}$, $H^2 = H_{\mu\nu\lambda}H^{\mu\nu\lambda}$ involve the totally antisymmetric field strength $H_{\mu\nu\lambda}$, defined in terms of the potential $B_{\mu\nu}$ as

$$H_{\mu\nu\rho} = \partial_\mu B_{\nu\rho} + \partial_\rho B_{\mu\nu} + \partial_\nu B_{\rho\mu}. \tag{5}$$

Indeed these equations are the Einstein field equations coupled with matters ϕ and H (see for instance [8]). Note that for the study of 4+1 dimensional homogeneous string cosmological models, we first consider it in $d+1$-dimension . We know that with the assumption of spatial homogeneity, the d-dimensional spatial submanifold is invariant under the action of a d-parametric isometry group G, and the metric of the space-time $G_{\mu\nu}$ can be factorized in a synchronous frame as [13]:

$$ds^2 = G_{\mu\nu}dx^\mu dx^\nu = -dt^2 + e_\alpha{}^i(x)g_{ij}(t)e_\beta{}^j(x)dx^\alpha dx^\beta, \tag{6}$$

where $\alpha, \beta = 1, ..., d$ are world indices in the spatial submanifold and i, j are indices for the bases of the isometry group G. Notice that vielbeins $e_\alpha{}^i(x)$ are dependent only on spatial coordinates [13]. Now, if we consider additional assumption that the dilaton field be space independent and the torsion field $H_{\mu\nu\rho}$ vanish then the equations(2)-(4) can be reduce to ordinary time differential equations for the variables $g_{ij}(t)$ and $\phi(t)$ [9]. Furthermore if we restrict our analysis to an anisotropic but diagonal matrix form for the invariant metric $g_{ij}(t)$ [1],

$$g_{ij}(t) = a_i^2(t)\delta_{ij}, \tag{7}$$

where $a_i(t)$ are scale factors; then the (0,0) and (i,i) components of the equation (2) have the following form respectively [9]:

$$\sum_{i=1}^{d}(\dot{H}_i + H_i^2) + \ddot{\phi} = 0, \tag{8}$$

$$\dot{H}_i + H_i\sum_{k=1}^{d}H_k + H_i\dot{\phi} + V_i = 0, \tag{9}$$

where the dot stands for derivative with respect to t; furthermore (i,0) components give the constraints[2]

$$\sum_{k=1}^{d}C^k_{ki}(H_i - H_k) = 0. \tag{10}$$

On the other hand, $i \neq j$ components of equation (2) give a set of another constraint equations

$$R_{i,j} = 0, \qquad i \neq j, \tag{11}$$

---

[1] Note that one can also use other form of g_{ij}; for example for four dimensional Lie groups one can use results of Ref [14].

[2] Here there is no summation over i.

and the dilaton equation (4) has the following form:

$$-2\ddot{\phi} - \dot{\phi}^2 - 2\dot{\phi}\sum_{k=1}^{d} H_k - \sum_{k=1}^{d} V_k - (\sum_{k=1}^{d} H_k)^2 - \sum_{k=1}^{d} {H_k}^2 - 2\sum_{k=1}^{d} \dot{H}_k = 0. \tag{12}$$

Note that in the above equations $H_i = \frac{\dot{a}_i}{a_i}$ are Hubble coefficients, C_{ij}^k are structure constant of the isometry Lie algebras $\mathbf{g}$ and $V_k(a_i)$ are potential functions that depend on the type of Lie group. In this way instead of equations $(2-4)$ one can use equations $(8-12)$ and solve them to obtain homogeneous string cosmological models over space-times with isometry group G.

Now, for 4+1 dimensional models the isometry groups are four dimensional real Lie groups. There are various classifications for these Lie algebras [15, 16, 17, 18]. Here in this paper we use the Patera et [17] classification in the next section. Now, for the four dimensional spatial submanifold one can introduce mean radius a in term of scale factors a_i as follows:

$$a^4 = a_1 a_2 a_3 a_4, \tag{13}$$

then using equations (8),(9),(12) we have

$$\ddot{\phi} + \dot{\phi}^2 + 4\dot{\phi}\frac{\dot{a}}{a} = 0. \tag{14}$$

On the other hand, introducing new time coordinate τ [8]

$$d\tau = a^{-4} e^{-\phi} dt, \tag{15}$$

the equation (14) can be rewritten as

$$\phi'' = 0, \tag{16}$$

where prime stands for derivative with respect to τ. The general solution of the above equation is

$$\phi = N\tau, \quad or \quad e^{-\phi} = e^{-N\tau}, \tag{17}$$

where N is a constant. Furthermore, with this new time coordinate the equation (9) may be rewritten as

$$\ln(a_i^2 e^{\phi})'' + 2a^8 e^{2\phi} V_i = 0. \tag{18}$$

On the other hand, equation (12) can be reexpressed as the following initial value equations:

$$\sum_{i<j} \ln(a_i^2 e^{\phi})' \ln(a_j^2 e^{\phi})' + 2\sum_i V_i a^8 e^{2\phi} - 2\sum_i \frac{a_i{}'}{a_i}\phi' - 4\phi'^2 = 0. \tag{19}$$

Now, it is enough to solve equation (18) with the constraints (10) and (11)and initial values equations (19). In order to solve equations (18) we must obtain potentials $V_i(a_j)$. In the next section we obtain metrics over four dimensional Lie groups, then in section four we obtain these potentials and solve equations (18).

3 Four dimensional real Lie algebras and metrics over their Lie groups

There are several classifications for real four dimensional Lie algebras (for instance see [15], [16], [17], [18]). Because in most cosmological applications the Patera and et al classification has been used (see for instance [12, 14] and references), here we apply this classification (Table 1). Now, to obtain the metrics of 4+1 dimensional anisotropic homogenouse space-times according to (6), we must first calculate vielbeins ${e_\alpha}^i(x)$ for Lie groups of table 1. To this end, we use the following relation:

$$g^{-1}dg = {e_\alpha}^i(x) X_i dx^\alpha, \qquad g \in G. \tag{20}$$

With the following parameterizations for the real four dimensional Lie groups G:

$$g = e^{x_1 X_1} e^{x_2 X_2} e^{x_3 X_3} e^{x_4 X_4}, \tag{21}$$

where $\{X_i\}$ and $\{x_i\}$ are generators and coordinates of Lie group, respectively. Then in general, for left invariant Lie algebra valued one forms we have:

$$\begin{aligned} g^{-1}dg = {} & dx_1 e^{-x_4X_4} e^{-x_3X_3} (e^{-x_2X_2} X_1 e^{x_2X_2}) e^{x_3X_3} e^{x_4X_4} + dx_2 e^{-x_4X_4} e^{-x_3X_3} X_2 e^{x_3X_3} e^{x_4X_4} \\ & + dx_3 e^{-x_4X_4} X_3 e^{x_4X_4} + dx_4 X_4. \end{aligned} \tag{22}$$

Now, we need to calculate expressions such as $e^{-x_iX_i} X_j e^{x_iX_i}$. Indeed, in[19] we have shown that:

$$e^{-x_iX_i} X_j e^{x_iX_i} = {(e^{x_i\mathcal{X}_i})_j}^k X_k, \tag{23}$$

where summation over index k is assumed and there is no summation over index i. In this formula ${(\mathcal{X}_i)_l}^j = -{f_{il}}^j$ are the adjoint representation of the Lie algebra. In this way one can calculate all left invariant one forms, which are shown in Table 1.

Table 1: 4-dimensional real Lie algebras, their isometry Lie algebras and left invariant one forms. [3]

Algebra	Non-zero commutation relations	Isometry	$g^{-1}dg$
$4A_1$	$[X_i, X_j] = 0$	$I \oplus R$	$dx^i X_i$
$A_2 + 2A_1$	$[X_1, X_2] = -X_2 - X_3, [X_3, X_1] = X_2 + X_3$	X_1, X_4	$dx_1[X_1 - (x_2 + x_3)(X_2 + X_3)] + dx_2 X_2 + dx_3(X_3 + x_4 X_4) + dx_4 X_4$
$2A_2$	$[X_1, X_2] = X_2, [X_3, X_4] = X_4$	X_1, X_3	$dx_1(X_1 + x_2 X_2) + dx_2 X_2 + dx_3(X_3 + x_4 X_4) + dx_4 X_4$
$II \oplus R$	$[X_2, X_3] = X_1$	X_1, X_2, X_4	$dx_1 X_1 + dx_2(X_2 + x_3 X_1) + dx_3 X_3 + dx_4 X_4$
$IV \oplus R$	$[X_1, X_2] = -X_2 + X_3, [X_1, X_3] = -X_3$	X_1, X_4	$dx_1[X_1 - x_2 X_2 + (x_2 - x_3)X_3] + dx_2 X_2 + dx_3 X_3 + dx_4 X_4$
$V \oplus R$	$[X_1, X_2] = -X_2, [X_1, X_3] = -X_3$	X_1, X_4	$dx_1\,(X_1 - x_2 X_2 - x_3 X_3) + dx_2 X_2 + dx_3 X_3 + dx_4 X_4$
$VI_0 \oplus R$	$[X_1, X_3] = X_2, [X_2, X_3] = X_1$	X_1, X_2, X_4	$dx_1(X_1 \cosh x_3 + X_2 \sinh x_3) + dx_2(X_2 \cosh x_3 + X_1 \sinh x_3) + dx_3 X_3 + dx_4 X_4$
$VI_a \oplus R$	$[X_1, X_2] = -aX_2 - X_3, [X_3, X_1] = X_2 + aX_3$	X_1, X_4	$dx_1[X_1 - (ax_2 + x_3)X_2 - (ax_3 + x_2)X_3] + dx_2 X_2 + dx_3 X_3 + dx_4 X_4$
$VII_0 \oplus R$	$[X_1, X_3] = -X_2, [X_2, X_3] = X_1$	X_1, X_2, X_4	$dx_1(X_1 \cos x_3 - X_2 \sin x_3) + dx_2(X_2 \cos x_3 + X_1 \sin x_3) + dx_3 X_3 + dx_4 X_4$
$VII_a \oplus R$	$[X_1, X_2] = -aX_2 + X_3, [X_3, X_1] = X_2 + aX_3$	X_1, X_4	$dx_1[X_1 - (ax_2 + x_3)X_2 - (-ax_3 + x_2)X_3] + dx_2 X_2 + dx_3 X_3 + dx_4 X_4$
$VIII \oplus R$	$[X_1, X_3] = -X_2, [X_2, X_3] = X_1, [X_1, X_2] = -X_3$	X_1, X_4	$dx_1(X_1 \cosh x_2 \cos x_3 - X_2 \cosh x_2 \sin x_3 - X_3 \sinh x_2) + dx_2(X_2 \cos x_3 + X_1 \sin x_3) + dx_3 X_3 + dx_4 X_4$
$IX \oplus R$	$[X_1, X_3] = -X_2, [X_2, X_3] = X_1, [X_1, X_2] = X_3$	X_1, X_4	$dx_1(X_1 \cos x_2 \cos x_3 - X_2 \cos x_2 \sin x_3 + X_3 \sin x_2) + dx_2(X_2 \cos x_3 + X_1 \sin x_3) + dx_3 X_3 + dx_4 X_4$
$A_{4,1}$	$[X_2, X_4] = X_1, [X_3, X_4] = X_2$	X_1, X_2, X_3	$dx_1 X_1 + dx_2(X_1 x_4 + X_2) + dx_3(X_3 + \frac{x_4^2}{2} X_1 + X_2 x_4) + dx_4 X_4$
$A^a{}_{4,2}$	$[X_1, X_4] = aX_1, [X_2, X_4] = X_2, [X_3, X_4] = X_2 + X_3$	X_1, X_2, X_3	$dx_1 X_1 e^{ax_4} + dx_2 X_2 e^{x_4} + dx_3(X_3 + X_2)e^{x_4} + dx_4 X_4$
$A^1{}_{4,2}$	$[X_1, X_4] = aX_1, [X_1, X_4] = X_1, [X_2, X_4] = X_2, [X_3, X_4] = X_2 + X_3$	X_1, X_2, X_3	$dx_1 X_1 e^{x_4} + dx_2 X_2 e^{x_4} + dx_3(X_3 + X_2)e^{x_4} + dx_4 X_4$
$A_{4,3}$	$[X_1, X_4] = X_1, [X_3, X_4] = X_2$	X_1, X_2, X_3	$dx_1 X_1 e^{x_4} + dx_2 X_2 + dx_3(X_3 + x_4 X_2) + dx_4 X_4$
$A_{4,4}$	$[X_1, X_4] = X_1, [X_2, X_4] = X_1 + X_2, [X_3, X_4] = X_2 + X_3$	X_1, X_2, X_3	$dx_1 X_1 e^{x_4} + dx_2(X_2 + x_4 X_1)e^{x_4} + dx_3(X_3 + X_2 x_4 + X_1 x_4^2/2)e^{x_4} + dx_4 X_4$
$A^{a,b}{}_{4,5}$	$[X_1, X_4] = X_1, [X_2, X_4] = aX_2, [X_3, X_4] = bX_3$	X_1, X_2, X_3	$dx_1 X_1 e^{x_4} + dx_2 X_2 e^{ax_4} + dx_3 X_3 e^{bx_4} + dx_4 X_4$
$A^{a,a}{}_{4,5}$	$[X_1, X_4] = X_1, [X_2, X_4] = aX_2, [X_3, X_4] = aX_3$	X_1, X_2, X_3	$dx_1 X_1 e^{x_4} + dx_2 X_2 e^{ax_4} + dx_3 X_3 e^{ax_4} + dx_4 X_4$
$A^{a,1}{}_{4,5}$	$[X_1, X_4] = X_1, [X_2, X_4] = aX_2, [X_3, X_4] = X_3$	X_1, X_2, X_3	$dx_1 X_1 e^{x_4} + dx_2 X_2 e^{ax_4} + dx_3 X_3 e^{x_4} + dx_4 X_4$
$A^{1,1}{}_{4,5}$	$[X_1, X_4] = X_1, [X_2, X_4] = X_2, [X_3, X_4] = X_3$	X_1, X_2, X_3	$dx_1 X_1 e^{x_4} + dx_2 X_2 e^{x_4} + dx_3 X_3 e^{x_4} + dx_4 X_4$
$A^{a,b}{}_{4,6}$	$[X_1, X_4] = aX_1, [X_2, X_4] = 2X_2 - X_3, [X_3, X_4] = X_2 + bX_3$	X_1, X_2, X_3	$dx_1 X_1 e^{ax_4} + e^{bx_4}[dx_2(X_2 \cos x_4 - X_3 \sin x_4) + dx_3(X_3 \cos x_4 + X_2 \sin x_4)] + dx_4 X_4$

---

[3] Note that for Lie algebras written as direct sum of three dimensional Lie algebras with one dimensional Lie algebras we write their isomorphism with Bianchi $\oplus R$ Lie algebras.

Table 1 : Continue

Algebra	Non-zero commutation relations	Isometry	$g^{-1}dg$
$A_{4,7}$	$[X_1,X_4]=2X_1,[X_2,X_4]=X_2,$ $[X_3,X_4]=X_2+X_3,[X_2,X_3]=X_1$	X_1,X_2	$dx_1X_1e^{2x_4}+dx_2(X_2e^{x_4}-x_3X_1e^{2x_4})+dx_3e^{x_4}(x_4X_2$ $+X_3)+dx_4X_4$
$A_{4,8}$	$[X_2,X_4]=X_2,[X_3,X_4]=-X_3,[X_2,X_3]=X_1$	X_1,X_2	$dx_1X_1+dx_2(X_2e^{x_4}+x_3X_1)+dx_3e^{-x_4}X_3+dx_4X_4$
$A^b_{4,9}$	$[X_1,X_4]=(1+b)X_1,[X_2,X_4]=X_2,$ $[X_3,X_4]=bX_3,[X_2,X_3]=X_1$	X_1,X_2	$dx_1X_1e^{(1+b)x_4}+dx_2(X_2e^{x_4}+x_3e^{(1+b)x_4}X_1)$ $+dx_3e^{x_4}(x_4X_2+X_3)+dx_4X_4$
$A^1_{4,9}$	$[X_1,X_4]=2X_1,[X_2,X_4]=X_2,[X_3,X_4]=X_3,$ $[X_2,X_3]=X_1$	X_1,X_2	$dx_1X_1e^{2x_4}+dx_2(X_2e^{x_4}+x_3e^{2x_4}X_1)+dx_3e^{x_4}X_3+dx_4X_4$
$A^0{}_{4,9}$	$[X_1,X_4]=X_1,[X_2,X_4]=X_2,[X_2,X_3]=X_1$	X_1,X_2	$dx_1X_1e^{x_4}+dx_2(X_2e^{x_4}+x_3e^{x_4}X_1)+dx_3X_3+dx_4X_4$
$A_{4,10}$	$[X_2,X_4]=-X_3,\ [X_3,X_4]=X_2,;[X_2,X_3]=X_1$	X_1,X_2	$dx_1X_1+dx_2(x_3X_1+X_2\cos x_4-X_3\sin x_4)+dx_3(X_3\cos x_4$ $+X_2\sin x_4)+dx_4X_4$
$A^a{}_{4,11}$	$[X_1,X_4]=2aX_1,[X_2,X_4]=aX_2-X_3,$ $[X_3,X_4]=X_2+aX_3,[X_2,X_3]=X_1$	X_1,X_2	$dx_1X_1e^{2ax_4}+dx_2(x_3e^{2ax_4}X_1+X_2e^{ax_4}\cos x_4-$ $X_3e^{ax_4}\sin x_4)+dx_3(X_3e^{ax_4}\cos x_4+X_2e^{ax_4}\sin x_4)+dx_4X_4$
$A_{4,12}$	$[X_1,X_4]=-X_2,[X_1,X_3]=X_1,[X_2,X_4]=X_1,$ $[X_2,X_3]=X_2$	X_1,X_2	$dx_1e^{x_3}(X_1\cos x_4-X_2\sin x_4)+dx_2e^{x_3}(X_2\cos x_4+X_1\sin x_4)$ $+dx_3X_3+dx_4X_4$

4 Exact solutions

Here, we construct the (i,i) component of equation (2) for metrics over all real four dimensional Lie groups of the table (1), then by comparing these equations with (9) we obtain potentials V_i, furthermore we try to solve equations (18) and obtain initial value equation (19).

4.1 Lie algebras $I \oplus R$

We obtain the potentials V_i as follows:

$$V_1 = V_2 = V_3 = V_4 = 0, \tag{24}$$

such that there are no constraint equations. Equations (18) have the following form:

$$\ln(a_i^2 e^{\phi})'' = 0, \quad i = 1, 2, ..., 4 \tag{25}$$

which admit the following general solution:

$$a_i^2 e^{\phi} = L_i e^{p_i \tau}, \tag{26}$$

where p_i and $L_i (i = 1, 2, ..., 4)$ are constants; furthermore from the initial value equation (19) we have the following restriction on $p_i s$:

$$\sum_{i<j} p_i p_j = N \sum_i p_i. \tag{27}$$

These results are extension of the results of [8].

4.2 Lie algebras $III \oplus R$

We obtain the potentials V_i as follows:

$$V_1 = -\frac{a_2^2}{2a_1^2a_3^2} - \frac{a_3^2}{2a_1^2a_2^2} - \frac{3}{a_1^2}, \;\; V_2 = \frac{a_2^2}{2a_1^2a_3^2} - \frac{a_3^2}{2a_1^2a_2^2} - \frac{2}{a_1^2},$$
$$V_3 = -\frac{a_2^2}{2a_1^2a_3^2} + \frac{a_3^2}{2a_1^2a_2^2} - \frac{2}{a_1^2}, \;\; V_4 = 0. \tag{28}$$

In this case the constraints imposed by (10) and (11) are

$$\begin{aligned} &a_2a_3 = Ba_1^2, \\ &a_2^2 + a_3^2 = 0, \end{aligned} \tag{29}$$

where B is a constant. The above equations are inconsistent; and there is no solution for this example.

4.3 Lie algebras $A_2 \oplus A_2$

We obtain the potentials V_i as follows:

$$V_1 = V_2 = \frac{1}{a_1^2}, \;\; V_3 = V_4 = \frac{1}{a_3^2}. \tag{30}$$

The constraint equations (10) impose in this case the restriction

$$\frac{a_1}{a_2} = B_1, \; \frac{a_3}{a_4} = B_2, \tag{31}$$

where B_1 and B_2 are constants. Equations (18) have the following forms

$$\begin{aligned} &\ln(a_1^2e^{\phi})'' - 2a_2^2a_3^2a_4^2e^{2\phi} = 0, \\ &\ln(a_2^2e^{\phi})'' - 2a_2^2a_3^2a_4^2e^{2\phi} = 0, \\ &\ln(a_3^2e^{\phi})'' - 2a_1^2a_2^2a_4^2e^{2\phi} = 0, \\ &\ln(a_4^2e^{\phi})'' - 2a_1^2a_2^2a_4^2e^{2\phi} = 0, \end{aligned} \tag{32}$$

which admit the following general solution:

$$\begin{aligned} a_1^2e^{\phi} = a_3^2e^{\phi} &= p_4^2\left(\frac{(B_1B_2)^2e^{N\tau}}{3\cosh^2(p_4\tau)}\right)^{\frac{1}{3}}, \\ a_2^2e^{\phi} &= p_4^2\left(\frac{(\frac{B_2}{B_1^2})^2e^{N\tau}}{3\cosh^2(p_4\tau)}\right)^{\frac{1}{3}}, \\ a_4^2e^{\phi} &= p_4^2\left(\frac{(\frac{B_1}{B_2^2})^2e^{N\tau}}{3\cosh^2(p_4\tau)}\right)^{\frac{1}{3}}, \end{aligned} \tag{33}$$

where p_4 is constant. From the initial value equation (19) we have the following restriction on p_4:

$$N^2 - 4B_1^2B_2^2p_4^6 = 0. \tag{34}$$

4.4 Lie algebras $II \oplus R$

We obtain the potentials V_i as follows:

$$V_1 = -V_2 = -V_3 = \frac{a_1^2}{2a_2^2 a_3^2}, \;\; V_4 = 0, \tag{35}$$

such that there are no constraint equations. Equations (18) have the following form:

$$\begin{aligned}
&\ln(a_1^2 e^{\phi})'' + a_1^4 a_4^2 e^{2\phi} = 0,\\
&\ln(a_2^2 e^{\phi})'' - a_1^4 a_4^2 e^{2\phi} = 0,\\
&\ln(a_3^2 e^{\phi})'' - a_1^4 a_4^2 e^{2\phi} = 0,\\
&\ln(a_4^2 e^{\phi})'' = 0,
\end{aligned} \tag{36}$$

which admit the following general solution:

$$\begin{aligned}
a_1^2 e^{\phi} &= \frac{p_1^2 \, e^{(\frac{N-p_4}{2})\tau}}{\sqrt{L_1}\cosh(p_1\tau)},\\
a_2^2 e^{\phi} &= L_2^2 \cosh(p_1\tau) e^{p_2\tau},\\
a_3^2 e^{\phi} &= L_3^2 \cosh(p_1\tau) e^{p_3\tau},\\
a_4^2 e^{\phi} &= L_1 e^{p_4\tau},
\end{aligned} \tag{37}$$

where $p_1, ..., p_4, L_1, L_2$ and L_3 are constants. Note that these solutions are extension of solutions [8] for Bianchi algebras II. From the initial value equation (19) we have the following restriction on $p_i s$:

$$N(p_2 + p_3 + N) + p_4(p_2 + p_3 + p_4) - 2p_2 p_3 + 2p_1^2 = 0. \tag{38}$$

4.5 Lie algebras $(IV \oplus R)$

We obtain the potentials V_i as follows:

$$V_1 = V_2 = -\frac{a_3^2}{2a_1^2 a_2^2} - \frac{2}{a_1^2}, \;\; V_3 = \frac{a_3^2}{2a_1^2 a_2^2} - \frac{2}{a_1^2}, \;\; V_4 = 0. \tag{39}$$

In this case the constraints imposed by (10) and (11) are as follows:

$$\begin{aligned}
&a_2 a_3 = B a_1^2,\\
&a_3 = 0,
\end{aligned} \tag{40}$$

where B is constant. There is no solution for these inconsistent constraints.

4.6 Lie algebras $A_{3,3} \oplus A_1 \approx V \oplus R$

In this case we obtain the potentials V_i as follows:

$$V_1 = V_2 = V_3 = \frac{2}{a_1^2}, \;\; V_4 = 0, \tag{41}$$

and the constraint imposed by (10) and (11) is

$$a_2 a_3 = B a_1^2. \tag{42}$$

Equations (18) have the following forms:

$$\begin{aligned}
&\ln(a_1^2 e^{\phi})'' - 4a_2^2 a_3^2 a_4^2 e^{2\phi} = 0,\\
&\ln(a_2^2 e^{\phi})'' - 4a_2^2 a_3^2 a_4^2 e^{2\phi} = 0,\\
&\ln(a_3^2 e^{\phi})'' - 4a_2^2 a_3^2 a_4^2 e^{2\phi} = 0,\\
&\ln(a_4^2 e^{\phi})'' = 0,
\end{aligned} \tag{43}$$

which admit the following general solution:

$$\begin{aligned}
a_1^2 e^{\phi} &= \frac{p_1\, e^{(\frac{N-p_4}{2})\tau}}{2B\sqrt{L_1}\sinh(p_1\tau)},\\
a_2^2 e^{\phi} &= \frac{L_2\, e^{p_2\tau}}{2\sinh(p_1\tau)},\\
a_3^2 e^{\phi} &= \frac{p_1^2\, e^{(N-p_2-p_4)\tau}}{2L_1 L_2\sinh(p_1\tau)},\\
a_4^2 e^{\phi} &= L_1 e^{p_4\tau},
\end{aligned} \tag{44}$$

where p_1, p_2, p_4, L_1 and L_2 are constants. From the initial value equation (19) we have the following restriction on $p_i s$:

$$N(p_2 + p_4 - N) - p_2(p_2 + p_4) - \left(p_4^2 - 3p_1^2\right) = 0. \tag{45}$$

4.7 Lie Algebras $A_{3,4} \oplus A_1 \approx VI_0 \oplus R$

We obtain the potentials V_i as follows:

$$V_1 = -V_2 = \frac{a_1^2}{2a_2^2 a_3^2} - \frac{a_2^2}{2a_1^2 a_3^2}, \;\; V_3 = -\frac{a_1^2}{2a_2^2 a_3^2} - \frac{a_2^2}{2a_1^2 a_3^2} - \frac{1}{a_3^2}, \;\; V_4 = 0, \tag{46}$$

such that there are no constraint equations. Equations (18) have the following forms:

$$\begin{aligned}
&\ln(a_1^2 e^{\phi})'' + (a_1^4 - a_2^4) a_4^2 e^{2\phi} = 0,\\
&\ln(a_2^2 e^{\phi})'' - (a_1^4 - a_2^4) a_4^2 e^{2\phi} = 0,\\
&\ln(a_3^2 e^{\phi})'' - (a_1^4 + a_2^4) a_4^2 e^{2\phi} - 2a_1^2 a_2^2 a_4^2 e^{2\phi} = 0,\\
&\ln(a_4^2 e^{\phi})'' = 0,
\end{aligned} \tag{47}$$

which admit the following general solution:

$$
\begin{aligned}
&a_1^2e^{\phi} = a_2^2e^{\phi} = \sqrt{L_1}\, e^{\frac{p_1}{2}\tau},\\
&a_3^2e^{\phi} = exp\left(\frac{4L_1L_2}{(p_1+p_4-N)^2}\, e^{(p_1+p_4-N)\tau}\right),\\
&a_4^2e^{\phi} = L_2e^{p_4\tau},
\end{aligned}
\tag{48}
$$

where L_1, L_2, p_1 and p_4 are constants. From the initial value equation (19) we have the following restriction on p_1 and p_4:

$$
p_1(p_1+4p_4-4N) - 4p_4N = 0. \tag{49}
$$

Not that it has not presented solution for Bianchi algebra VI_0 in [8]

4.8 Lie Algebras $VI_b \oplus R$

We obtain the potentials V_i as follows:

$$
\begin{aligned}
&V_1 = -\frac{a_2^2}{2a_1^2a_3^2} - \frac{a_3^2}{2a_1^2a_2^2} - \frac{1+2b^2}{a_1^2},\ \ V_2 = \frac{a_2^2}{2a_1^2a_3^2} - \frac{a_3^2}{2a_1^2a_2^2} - \frac{2b^2}{a_1^2},\\
&V_3 = -\frac{a_2^2}{2a_1^2a_3^2} + \frac{a_3^2}{2a_1^2a_2^2} - \frac{2b^2}{a_1^2},\ \ V_4 = 0.
\end{aligned}
\tag{50}
$$

In this case, the constraints imposed by (10) and (11) are

$$
\begin{aligned}
&a_2a_3 = Ba_1^2,\\
&a_2^2 + a_3^2 = 0,
\end{aligned}
\tag{51}
$$

where B is constant; so that, within the present context, all solutions are singular.

4.9 Lie algebras $VII_0 \oplus R$

We obtain the potentials V_i as follows:

$$
V_1 = -V_2 = \frac{a_1^2}{2a_2^2a_3^2} - \frac{a_2^2}{2a_1^2a_3^2},\ \ V_3 = -\frac{a_1^2}{2a_2^2a_3^2} - \frac{a_2^2}{2a_1^2a_3^2} + \frac{1}{a_3^2},\ \ V_4 = 0, \tag{52}
$$

such that there are no constraint equations. Equations (18) have the following forms:

$$
\begin{aligned}
&\ln(a_1^2e^{\phi})'' + (a_1^4 - a_2^4)a_4^2e^{2\phi} = 0,\\
&\ln(a_2^2e^{\phi})'' - (a_1^4 - a_2^4)a_4^2e^{2\phi} = 0,\\
&\ln(a_3^2e^{\phi})'' - (a_1^4 + a_2^4)a_4^2e^{2\phi} + 2a_1^2a_2^2a_4^2e^{2\phi} = 0,\\
&\ln(a_4^2e^{\phi})'' = 0,
\end{aligned}
\tag{53}
$$

which admit the following general solution

$$\begin{aligned}
&a_1^2 e^{\phi} = a_2^2 e^{\phi} = \sqrt{L_1}\, e^{\frac{p_1}{2}\tau},\\
&a_3^2 e^{\phi} = L_2 e^{p_3\tau},\\
&a_4^2 e^{\phi} = L_3 e^{p_4\tau},
\end{aligned} \tag{54}$$

where L_1, L_2, L_3, p_1, p_3 and p_4 are constants. Furthermore from the initial value equation (19) we have the following restriction on $p_i s$:

$$\frac{p_1^2}{4} + p_1 p_3 + p_1 p_4 + p_3 p_4 - (p_1 + p_3 + p_4)\, N = 0. \tag{55}$$

Note that this solution and initial value equation are the same as for the Lie algebra $I \oplus R$.

4.10 Lie algebras $VII_b \oplus R$

We obtain the potentials V_i as follows:

$$\begin{aligned}
&V_1 = -\frac{a_3^2}{2a_1^2 a_2^2} - \frac{a_2^2}{2a_1^2 a_3^2} - \frac{2b^2 - 1}{a_1^2}, \;\; V_2 = \frac{a_2^2}{2a_1^2 a_3^2} - \frac{a_3^2}{2a_1^2 a_2^2} - \frac{2b^2}{a_1^2},\\
&V_3 = -\frac{a_2^2}{2a_1^2 a_3^2} + \frac{a_3^2}{2a_1^2 a_2^2} - \frac{2b^2}{2a_1^2}, \;\; V_4 = 0.
\end{aligned} \tag{56}$$

In this case the constraints imposed by (10), (11) are

$$\begin{aligned}
&a_2 a_3 = B a_1^2,\\
&a_2^2 - a_3^2 = 0,
\end{aligned} \tag{57}$$

where B is constant. Equations (18) have the following forms:

$$\begin{aligned}
&\ln(a_1^2 e^{\phi})'' - \left((a_2^2 - a_3^2)^2 + 4b^2 a_2^2 a_3^2\right) a_4^2 e^{2\phi} = 0,\\
&\ln(a_2^2 e^{\phi})'' - 4b^2 a_2^2 a_3^2 a_4^2 e^{2\phi} = 0,\\
&\ln(a_3^2 e^{\phi})'' - 4b^2 a_2^2 a_3^2 a_4^2 e^{2\phi} = 0,\\
&\ln(a_4^2 e^{\phi})'' = 0,
\end{aligned} \tag{58}$$

which admit the following general solution:

$$\begin{aligned}
&a_1^2 e^{\phi} = \frac{p_1\, e^{(\frac{N - p_4}{2})\tau}}{2bB\sqrt{L_1}\sinh(p_1\tau)},\\
&a_2^2 e^{\phi} = a_3^2 e^{\phi} = \frac{p_1\, e^{(\frac{N - p_4}{2})\tau}}{2b\sqrt{L_1}\sinh(p_1\tau)},\\
&a_4^2 e^{\phi} = L_1 e^{p_4\tau},
\end{aligned} \tag{59}$$

where L_1, p_1 and p_4 are constants. From the initial value equation (19) we have the following restriction on p_1 and p_4:

$$4p_1^2 - p_4^2 - N^2 + \frac{2}{3} N p_4 = 0. \tag{60}$$

4.11 Lie algebras $VIII \oplus R$

We obtain the potentials V_i as follows:

$$V_1 = \frac{a_1^2}{2a_2^2a_3^2} - \frac{a_2^2}{2a_1^2a_3^2} - \frac{2a_2^2 + a_3^2}{2a_1^2a_2^2}, \;\; V_2 = -\frac{a_1^2}{2a_2^2a_3^2} + \frac{a_2^2}{2a_1^2a_3^2} - \frac{2a_1^2 + a_3^2}{2a_1^2a_2^2},$$
$$V_3 = -\frac{a_2^2}{2a_1^2a_3^2} + \frac{a_3^2}{2a_1^2a_2^2} - \frac{a_1^2 - 2a_2^2}{2a_2^2a_3^2}, \;\; V_4 = 0, \tag{61}$$

such that there are no constraint equations. Equations (18) have the following forms:

$$\begin{aligned}
&\ln(a_1^2e^{\phi})'' + \left(a_1^4 - (a_2^2 + a_3^2)^2\right) a_4^2e^{2\phi} = 0,\\
&\ln(a_2^2e^{\phi})'' + \left(a_2^4 - (a_1^2 + a_3^2)^2\right) a_4^2e^{2\phi} = 0,\\
&\ln(a_3^2e^{\phi})'' + \left(a_3^4 - (a_1^2 - a_2^2)^2\right) a_4^2e^{2\phi} = 0,\\
&\ln(a_4^2e^{\phi})'' = 0,
\end{aligned} \tag{62}$$

which admit the following general solution:

$$\begin{aligned}
&a_1^2e^{\phi} = a_2^2e^{\phi} = \frac{p_1^2\, e^{(\frac{N-p_4}{2})\tau} \cosh(p_3\tau)}{\sqrt{L_1}p_3 \sinh^2(p_1\tau)},\\
&a_3^2e^{\phi} = \frac{p_3\, e^{(\frac{N-p_4}{2})\tau}}{\sqrt{L_1}\cosh(p_3\tau)},\\
&a_4^2e^{\phi} = L_1e^{p_4\tau},
\end{aligned} \tag{63}$$

where L_1, p_1, p_3 and p_4 are constants. From the initial value equation (19) we have the following restriction on p_is:

$$4p_1^2 - p_3^2 - \frac{3}{4}p_4^2 + \frac{5}{4}Np_4 - \frac{3}{4}N^2 = 0. \tag{64}$$

4.12 Lie algebras $IX \oplus R$

We obtain the potentials V_i as follows:

$$V_1 = \frac{a_1^2}{2a_2^2a_3^2} - \frac{a_2^2}{2a_1^2a_3^2} + \frac{2a_2^2 - a_3^2}{2a_1^2a_2^2}, \;\; V_2 = -\frac{a_1^2}{2a_2^2a_3^2} + \frac{a_2^2}{2a_1^2a_3^2} + \frac{2a_1^2 - a_3^2}{2a_1^2a_2^2},$$
$$V_3 = -\frac{a_2^2}{2a_1^2a_3^2} + \frac{a_3^2}{2a_1^2a_2^2} - \frac{a_1^2 - 2a_2^2}{2a_2^2a_3^2}, \;\; V_4 = 0, \tag{65}$$

such that there are no constraint equations. Equations (18) have the following forms:

$$\begin{aligned}
&\ln(a_1^2e^{\phi})'' + \left(a_1^4 - (a_2^2 - a_3^2)^2\right) a_4^2e^{2\phi} = 0,\\
&\ln(a_2^2e^{\phi})'' + \left(a_2^4 - (a_1^2 - a_3^2)^2\right) a_4^2e^{2\phi} = 0,\\
&\ln(a_3^2e^{\phi})'' + \left(a_3^4 - (a_1^2 - a_2^2)^2\right) a_4^2e^{2\phi} = 0,\\
&\ln(a_4^2e^{\phi})'' = 0,
\end{aligned} \tag{66}$$

which admit the following general solution:

$$\begin{aligned}
&a_1^2 e^{\phi} = a_2^2 e^{\phi} = \frac{p_1^2\, e^{(\frac{N-p_4}{2})\tau}\cosh(p_3\tau)}{\sqrt{L_1}p_3\cosh^2(p_1\tau)},\\
&a_3^2 e^{\phi} = \frac{p_3\, e^{(\frac{N-p_4}{2})\tau}}{\sqrt{L_1}\cosh(p_3\tau)},\\
&a_4^2 e^{\phi} = L_1 e^{p_4\tau},
\end{aligned}\tag{67}$$

where L_1, p_1, p_3 and p_4 are constants. From the initial value equation (19) we have the following restriction on $p_i s$:

$$4p_1^2 - p_3^2 - \frac{3}{4}p_4^2 + \frac{5}{4}Np_4 - \frac{3}{4}N^2 = 0.\tag{68}$$

Note that equations (63) and (67) for $p_4 = 0$ are the same initial values equations of the Ref [8] for Lie algebras $VIII$ and IX respectively such that the scalar field play the role of antisymmetric field i.e. $\frac{3}{4}N^2 = A^2$; where A is a parameter of antisymmetric field [8].

4.13 Lie algebras $A_{4,1}$

We obtain the potentials V_i as follows:

$$V_1 = \frac{a_1^2}{2a_2^2a_4^2},\ \ V_2 = -\frac{a_1^2}{2a_2^2a_4^2} + \frac{a_2^2}{2a_3^2a_4^2},\ \ V_3 = -\frac{a_2^2}{2a_3^2a_4^2},\ \ V_4 = -\frac{a_1^2}{2a_2^2a_4^2} - \frac{a_2^2}{2a_3^2a_4^2},\tag{69}$$

such that there are no constraint equations. Equations (18) have the following forms:

$$\begin{aligned}
&\ln(a_1^2 e^{\phi})'' + a_1^4 a_3^2 e^{2\phi} = 0,\\
&\ln(a_2^2 e^{\phi})'' + \left(a_1^2 a_2^4 - a_3^2 a_1^4\right) e^{2\phi} = 0,\\
&\ln(a_3^2 e^{\phi})'' - a_1^2 a_2^4 e^{2\phi} = 0,\\
&\ln(a_4^2 e^{\phi})'' - \left(a_1^2 a_2^4 + a_3^2 a_1^4\right) e^{2\phi} = 0,
\end{aligned}\tag{70}$$

which admit the following general solution:

$$\begin{aligned}
&a_1^2 e^{\phi} = \frac{2p_2^2}{L_2^2\cosh^2(p_2\tau)}\, e^{(N-2p_2)\tau},\\
&a_2^2 e^{\phi} = L_2\, e^{p_2\tau},\\
&a_3^2 e^{\phi} = \frac{L_2^4\cosh^2(p_2\tau)}{2p_2^2}\, e^{(4p_2-N)\tau},\\
&a_4^2 e^{\phi} = L_4\cosh^4(p_2\tau) e^{p_4\tau},
\end{aligned}\tag{71}$$

where L_2, L_4, p_2 and p_4 are constants. From the initial value equation (19) we have the following restriction on p_2 and p_4:

$$N\left(N + p_4 - 3p_2\right) + 7p_2p_4 = 0.\tag{72}$$

4.14 Lie algebras $(A^b_{4,2})$

We obtain the potentials V_i as follows:

$$\begin{aligned}
&V_1 = -\frac{2+b^2}{a_4^2}, \;\; V_2 = \frac{a_2^2}{2a_3^2a_4^2} - \frac{2+b}{a_4^2}, \\
&V_3 = -\frac{a_2^2}{2a_3^2a_4^2} - \frac{2+b}{a_4^2}, \;\; V_4 = -\frac{a_2^2}{2a_3^2a_4^2} - \frac{2+b^2}{a_4^2}.
\end{aligned} \tag{73}$$

In this case the constraints imposed by (10) and (11) are

$$\begin{aligned}
&a_1^b a_2 a_3 = B a_4^{b+2}, \\
&a_2 = 0,
\end{aligned} \tag{74}$$

where B is constant; so that, within the present context, all solutions are singular.

4.15 Lie algebras $(A^1_{4,2})$

We obtain the potentials V_i as follows:

$$\begin{aligned}
&V_1 = -\frac{3}{a_4^2}, \;\; V_2 = \frac{a_2^2}{2a_3^2a_4^2} - \frac{3}{a_4^2}, \\
&V_3 = -\frac{a_2^2}{2a_3^2a_4^2} - \frac{3}{a_4^2}, \;\; V_4 = -\frac{a_2^2}{2a_3^2a_4^2} - \frac{3}{a_4^2}.
\end{aligned} \tag{75}$$

In this case the constraints imposed by (10) and (11) are

$$\begin{aligned}
&a_1 a_2 a_3 = B a_4^3, \\
&a_2 = 0,
\end{aligned} \tag{76}$$

where B is constant; so that, within the present context, all solutions are singular.

4.16 Lie algebras $(A_{4,3})$

We obtain the potentials V_i as follows:

$$V_1 = -\frac{1}{a_4^2}, \;\; V_2 = \frac{a_2^2}{2a_3^2a_4^2}, \;\; V_3 = -\frac{a_2^2}{2a_3^2a_4^2}, \;\; V_4 = -\frac{a_2^2}{2a_3^2a_4^2} - \frac{1}{a_4^2}. \tag{77}$$

In this case the constraints imposed by (10) and (11) are

$$\begin{aligned}
&a_1 = B a_4, \\
&a_2 = 0,
\end{aligned} \tag{78}$$

where B is constant; so that, within the present context, all solutions are singular.

4.17 Lie algebras $A_{4,4}$

We obtain the potentials V_i as follows:

$$
\begin{aligned}
&V_1 = \frac{a_1^2}{2a_2^2a_4^2} - \frac{3}{a_4^2},\ \ V_2 = -\frac{a_2^2}{2a_3^2a_4^2} - \frac{3}{a_4^2},\\
&V_3 = -\frac{a_1^2}{2a_2^2a_4^2} + \frac{a_2^2}{2a_3^2a_4^2} - \frac{3}{a_4^2},\ \ V_4 = -\frac{a_1^2}{2a_2^2a_4^2} - \frac{a_2^2}{2a_3^2a_4^2} - \frac{3}{a_4^2}.
\end{aligned}
\tag{79}
$$

In this case the constraints imposed by (10) and (11) are

$$
\begin{aligned}
&a_1a_2a_3 = Ba_4^3,\\
&a_1 = 0,
\end{aligned}
\tag{80}
$$

where B is constant; so that, within the present context, all solutions are singular.

4.18 Lie algebras $A_{4,5}^{(a,b)}$

We obtain the potentials V_i as follows:

$$
\begin{aligned}
&V_1 = -\frac{a+b+1}{a_4^2},\ \ V_2 = -\frac{a\,(a+b+1)}{a_4^2},\\
&V_3 = -\frac{b\,(a+b+1)}{a_4^2},\ \ V_4 = -\frac{a^2+b^2+1}{a_4^2}.
\end{aligned}
\tag{81}
$$

In this case the constraint equations (10) impose the following restriction:

$$
a_1a_2^aa_3^b = a_4^{a+b+1}.
\tag{82}
$$

Equations (18) have the following forms:

$$
\begin{aligned}
&\ln(a_1^2e^{\phi})'' - 2\,(a+b+1)\,a_1^2a_2^2a_3^2\,e^{2\phi} = 0,\\
&\ln(a_2^2e^{\phi})'' - 2a\,(a+b+1)\,a_1^2a_2^2a_3^2\,e^{2\phi} = 0,\\
&\ln(a_3^2e^{\phi})'' - 2b\,(a+b+1)\,a_1^2a_2^2a_3^2\,e^{2\phi} = 0,\\
&\ln(a_4^2e^{\phi})'' - 2\left(a^2+b^2+1\right)a_1^2a_2^2a_3^2\,e^{2\phi} = 0,
\end{aligned}
\tag{83}
$$

which for $a+b \neq -1$ admit the following general solution:

$$
\begin{aligned}
&a_1^2e^{\phi} = \frac{L_1^b(\frac{p_4}{a+b+1})^{\frac{2}{a+b+1}}\,e^{p_1\tau}}{\sinh(p_4\tau)^{\frac{2}{a+b+1}}},\\
&a_2^2e^{\phi} = \frac{L_2^{\frac{b}{a}}(\frac{p_4}{a+b+1})^{\frac{2a}{a+b+1}}\,e^{p_2\tau}}{\sinh(p_4\tau)^{\frac{2a}{a+b+1}}},\\
&a_3^2e^{\phi} = \frac{(\frac{p_4}{a+b+1})^{\frac{2b}{a+b+1}}\,e^{(N-p_1-p_2)\tau}}{L_1L_2\sinh(p_4\tau)^{\frac{2b}{a+b+1}}},\\
&a_4^2e^{\phi} = \frac{(\frac{p_4}{a+b+1})^{\frac{2(a^2+b^2+1)}{(a+b+1)^2}}\,e^{\left(\frac{p_2(a-b)+p_1(1-b)+bN}{a+b+1}\right)\tau}}{\sinh(p_4\tau)^{\frac{2\left(a^2+b^2+1\right)}{(a+b+1)^2}}},
\end{aligned}
\tag{84}
$$

where L_1, L_2, p_1, p_2 and p_4 are constants. From the initial value equation (19) we have the following restriction on $p_i s$:

$$p_1^2 + p_2^2 + N^2 + p_1 p_2 - N(p_1 + p_2) - 2\left(1 + \frac{a^2 + b^2 + 1}{(a + b + 1)^2}\right) p_4^2 = 0. \tag{85}$$

For $a + b = -1$ solution is similar to solution of VI_0 model.

4.19 Lie algebras $A_{4,5}^{(a,a)}$

We obtain the potentials V_i as follows:

$$V_1 = -\frac{2a + 1}{a_4^2}, \; V_2 = V_3 = aV_1 = -\frac{a(2a + 1)}{a_4^2}, \; V_4 = -\frac{2a^2 + 1}{a_4^2}. \tag{86}$$

The constraint equations (10) impose the following restriction:

$$a_1 a_2^a a_3^a = a_4^{2a+1}. \tag{87}$$

Equations (18) have the following forms:

$$\begin{aligned}
&\ln(a_1^2 e^{\phi})'' - 2\,(2a + 1)\, a_1^2 a_2^2 a_3^2\, e^{2\phi} = 0,\\
&\ln(a_2^2 e^{\phi})'' - 2a\,(2a + 1)\, a_1^2 a_2^2 a_3^2\, e^{2\phi} = 0,\\
&\ln(a_3^2 e^{\phi})'' - 2a\,(2a + 1)\, a_1^2 a_2^2 a_3^2\, e^{2\phi} = 0,\\
&\ln(a_4^2 e^{\phi})'' - 2\left(2a^2 + 1\right) a_1^2 a_2^2 a_3^2\, e^{2\phi} = 0,
\end{aligned} \tag{88}$$

which for $a \neq \frac{-1}{2}$ admit the following general solution:

$$\begin{aligned}
a_1^2 e^{\phi} &= \frac{L_1^a (\frac{p_4}{2a+1})^{\frac{2}{2a+1}}\; e^{p_1 \tau}}{\sinh(p_4 \tau)^{\frac{2}{2a+1}}},\\
a_2^2 e^{\phi} &= \frac{L_2 (\frac{p_4}{2a+1})^{\frac{2a}{2a+1}}\; e^{p_2 \tau}}{\sinh(p_4 \tau)^{\frac{2a}{2a+1}}},\\
a_3^2 e^{\phi} &= \frac{(\frac{p_4}{2a+1})^{\frac{2a}{2a+1}}\; e^{(N - p_1 - p_2)\tau}}{L_1 L_2 \sinh(p_4 \tau)^{\frac{2a}{2a+1}}},\\
a_4^2 e^{\phi} &= \frac{(\frac{p_4}{2a+1})^{\frac{2(2a^2+1)}{(2a+1)^2}}\; e^{\left(\frac{p_1(1-a) + aN}{2a+1}\right)\tau}}{\sinh(p_4 \tau)^{\frac{2\left(2a^2+1\right)}{(2a+1)^2}}},
\end{aligned} \tag{89}$$

where L_1, L_2, p_1, p_2 and p_4 are constants. From the initial value equation (19) we have the following restriction on $p_i s$:

$$p_1^2 + p_2^2 + N^2 + p_1 p_2 - N(p_1 + p_2) - 2\left(1 + \frac{2a^2 + 1}{(2a + 1)^2}\right) p_4^2 = 0. \tag{90}$$

For $a = \frac{-1}{2}$ solution is similar to solution of VI_0 model. Note that for this Lie algebra all equations, constraints solutions and initial value conditions are the same as for Lie algebra $A_{4,5}^{(a,b)}$ for $b = a$.

4.20 Lie algebras $A_{4,5}^{(a,1)}$

We obtain the potentials V_i as follows:

$$V_1 = V_3 = -\frac{a+2}{a_4^2}, \;\; V_2 = aV_1 = -\frac{a\,(a+2)}{a_4^2}, \;\; V_4 = -\frac{a^2+2}{a_4^2}. \tag{91}$$

The constraint equations (10) impose in this case the restriction

$$a_1 a_2^a a_3 = a_4^{a+2}. \tag{92}$$

Equations (18) have the following forms:

$$\begin{aligned}
&\ln(a_1^2 e^{\phi})'' - 2\,(a+2)\,a_1^2 a_2^2 a_3^2\, e^{2\phi} = 0,\\
&\ln(a_2^2 e^{\phi})'' - 2a\,(a+2)\,a_1^2 a_2^2 a_3^2\, e^{2\phi} = 0,\\
&\ln(a_3^2 e^{\phi})'' - 2\,(a+2)\,a_1^2 a_2^2 a_3^2\, e^{2\phi} = 0,\\
&\ln(a_4^2 e^{\phi})'' - 2\left(a^2+2\right) a_1^2 a_2^2 a_3^2\, e^{2\phi} = 0,
\end{aligned} \tag{93}$$

which admit the following general solution:

$$\begin{aligned}
a_1^2 e^{\phi} &= \frac{L_1^b (\frac{p_4}{a+2})^{\frac{2}{a+2}}\, e^{p_1 \tau}}{\sinh(p_4\tau)^{\frac{2}{a+2}}},\\
a_2^2 e^{\phi} &= \frac{L_2 (\frac{p_4}{a+2})^{\frac{2a}{a+2}}\, e^{p_2 \tau}}{\sinh(p_4\tau)^{\frac{2a}{a+2}}},\\
a_3^2 e^{\phi} &= \frac{(\frac{p_4}{a+2})^{\frac{2}{a+2}}\, e^{(N-p_1-p_2)\tau}}{(L_1 L_2)^a \sinh(p_4\tau)^{\frac{2}{a+2}}},\\
a_4^2 e^{\phi} &= \frac{(\frac{p_4}{a+2})^{\frac{2(a^2+2)}{(a+2)^2}}\, e^{\left(\frac{p_2(a-1)+N}{a+2}\right)\tau}}{\sinh(p_4\tau)^{\frac{2(a^2+2)}{(a+2)^2}}},
\end{aligned} \tag{94}$$

where L_1, L_2, p_1, p_2 and p_4 are constants. From the initial value equation (19) we have the following restriction on p_i:

$$p_1^2 + p_2^2 + N^2 + p_1 p_2 - N(p_1+p_2) - 2\left(1 + \frac{a^2+2}{(a+2)^2}\right) p_4^2 = 0. \tag{95}$$

Note that the results for this algebra are same as for Lie algebra $A_{4,5}^{(a,b)}$ for $b=1$.

4.21 Lie algebras $A_{4,5}^{(1,1)}$

We obtain the potentials V_i as follows:

$$V_1 = V_2 = V_3 = V_4 = -\frac{3}{a_4^2}. \tag{96}$$

The constraint equations (10) impose the following restriction:

$$a_1 a_2 a_3 = a_4^3. \tag{97}$$

Equations (18) have the following forms:

$$\begin{aligned}
&\ln(a_1^2 e^{\phi})'' - 6a_1^2 a_2^2 a_3^2\, e^{2\phi} = 0,\\
&\ln(a_2^2 e^{\phi})'' - 6a_1^2 a_2^2 a_3^2\, e^{2\phi} = 0,\\
&\ln(a_3^2 e^{\phi})'' - 6a_1^2 a_2^2 a_3^2\, e^{2\phi} = 0,\\
&\ln(a_4^2 e^{\phi})'' - 6a_1^2 a_2^2 a_3^2\, e^{2\phi} = 0,
\end{aligned} \tag{98}$$

which admit the following general solution:

$$\begin{aligned}
a_1^2 e^{\phi} &= \frac{L_1 (\frac{p_4}{3})^{\frac{2}{3}}\, e^{p_1 \tau}}{\sinh(p_4 \tau)^{\frac{2}{3}}},\\
a_2^2 e^{\phi} &= \frac{L_2 (\frac{p_4}{3})^{\frac{2}{3}}\, e^{p_2 \tau}}{\sinh(p_4 \tau)^{\frac{2}{3}}},\\
a_3^2 e^{\phi} &= \frac{(\frac{p_4}{3})^{\frac{2}{3}}\, e^{(N-p_1-p_2)\tau}}{(L_1 L_2)\sinh(p_4 \tau)^{\frac{2}{3}}},\\
a_4^2 e^{\phi} &= \frac{(\frac{p_4}{3})^{\frac{2}{3}}\, e^{\frac{N}{3}\tau}}{\sinh(p_4 \tau)^{\frac{2}{3}}},
\end{aligned} \tag{99}$$

where L_1, L_2, p_1, p_2 and p_4 are constants. From the initial value equation (19) we have the following restriction on p_i:

$$p_1^2 + p_2^2 + N^2 + p_1 p_2 - N(p_1 + p_2) - \frac{8}{3} p_4^2 = 0. \tag{100}$$

Note that the results for this algebra are same as for Lie algebra $A_{4,5}^{(a,1)}$ for $a = 1$.

4.22 Lie algebras $A_{4,6}^{(a,b)}$

We obtain the potentials V_i as follows:

$$\begin{aligned}
&V_1 = -\frac{a(a+2b)}{a_4^2},\;\; V_2 = \frac{a_2^2}{2a_3^2 a_4^2} - \frac{a_3^2}{2a_2^2 a_4^2} - \frac{b(a+2b)}{a_4^2},\\
&V_3 = -\frac{a_2^2}{2a_3^2 a_4^2} + \frac{a_3^2}{2a_2^2 a_4^2} - \frac{b(a+2b)}{a_4^2},\;\; V_4 = -\frac{a_2^2}{2a_3^2 a_4^2} - \frac{a_3^2}{2a_2^2 a_4^2} - \frac{(a^2+2b^2-1)}{a_4^2}.
\end{aligned} \tag{101}$$

The constraint equations (10) impose in this case the restriction

$$a_1^a (a_2 a_3)^b = B a_4^{a+2b}. \tag{102}$$

where B is a constant. Equations (18) have the following forms:

$$
\begin{aligned}
&\ln(a_1^2 e^{\phi})'' - 2\left(a^2+2ab\right) a_1^2 a_2^2 a_3^2 e^{2\phi} = 0,\\
&\ln(a_2^2 e^{\phi})'' - a_1^2\left(a_3^4 - a_2^4 + 2b(2b^2+a)a_2^2 a_3^2\right) e^{2\phi} = 0,\\
&\ln(a_3^2 e^{\phi})'' - a_1^2\left(a_2^4 - a_3^4 + 2b(2b^2+a)a_2^2 a_3^2\right) e^{2\phi} = 0,\\
&\ln(a_4^2 e^{\phi})'' - a_1^2\left(a_2^4 + a_3^4 + 2(2b^2+a^2-1)a_2^2 a_3^2\right) e^{2\phi} = 0,
\end{aligned}
\tag{103}
$$

which for $a \neq 2b$ admit the following general solution:

$$
\begin{aligned}
a_1^2 e^{\phi} &= \frac{L_1 p_1\, e^{p_1\tau}}{\sinh(p_1\tau)^{\frac{2a}{a+2b}}},\\
a_2^2 e^{\phi} = a_3^2 e^{\phi} &= \frac{\sqrt{p_1}\, e^{(\frac{N-p_1}{2})\tau}}{\sqrt{L_1}(a+2b)\sinh(p_1\tau)^{\frac{2b}{a+2b}}},\\
a_4^2 e^{\phi} &= \left(\frac{L_1^{a-b}}{B^2}\frac{p_1^{a+b}\, e^{((a-b)p_1+bN)\tau}}{\sinh(p_1\tau)^{2(\frac{a^2+2b^2}{a+2b})}}\right)^{\frac{1}{a+2b}},
\end{aligned}
\tag{104}
$$

where L_1 and p_1 are constants. From the initial value equation (19) we have the following restriction on p_1:

$$
(13a^2+36b^2+20ab)p_1^2 - (3a^2+4b^2+8ab)N^2 + 6a(a+2b)Np_1 = 0. \tag{105}
$$

But for the case $a = 2b$, its solution is similar to $VI_0 \oplus R$ model solution.

4.23 Lie algebras $A_{4,7}$

We obtain the potentials V_i as follows:

$$
\begin{aligned}
&V_1 = \frac{a_1^2}{2a_2^2a_3^2} - \frac{8}{a_4^2},\;\; V_2 = -\frac{a_1^2}{2a_2^2a_3^2} + \frac{a_2^2}{2a_3^2a_4^2} - \frac{4}{a_4^2},\\
&V_3 = -\frac{a_1^2}{2a_2^2a_3^2} - \frac{a_2^2}{2a_3^2a_4^2} - \frac{4}{a_4^2},\;\; V_4 = -\frac{a_2^2}{2a_3^2a_4^2} - \frac{6}{a_4^2}.
\end{aligned}
\tag{106}
$$

In this case the constraints imposed by (10) and (11) are

$$
\begin{aligned}
&a_1^2 a_2 a_3 = B a_4^4,\\
&a_2 = 0,
\end{aligned}
\tag{107}
$$

where B is constant; so that, within the present context, all solutions are singular.

4.24 Lie algebras $A_{4,8}$

We obtain the potentials V_i as follows:

$$
V_1 = \frac{a_1^2}{2a_2^2a_3^2},\;\; V_2 = -\frac{a_1^2}{2a_2^2a_3^2},\;\; V_3 = -\frac{a_1^2}{2a_2^2a_3^2},\;\; V_4 = -\frac{2}{a_4^2}. \tag{108}
$$

The constraint equations (10) impose in this case the restriction

$$a_2 = Ba_3, \tag{109}$$

where B is constant. In this case equations (18) have the following forms:

$$\begin{aligned}
&\ln(a_1^2 e^{\phi})'' + a_1^4 a_4^2 e^{2\phi} = 0,\\
&\ln(a_2^2 e^{\phi})'' - a_1^4 a_4^2 e^{2\phi} = 0,\\
&\ln(a_3^2 e^{\phi})'' - a_1^4 a_4^2 e^{2\phi} = 0,\\
&\ln(a_4^2 e^{\phi})'' - 4a_1^2 a_2^2 a_3^2 e^{2\phi} = 0,
\end{aligned} \tag{110}$$

which admit the following general solution:

$$\begin{aligned}
a_1^2 e^{\phi} &= \frac{3p_1\, e^{p_1\tau}}{2(L_1B)^2} \sinh(p_1\tau)^2,\\
a_2^2 e^{\phi} &= \frac{L_1 B^2 \sqrt{p_1}\, e^{(\frac{N-p_1}{2})\tau}}{\sinh(p_1\tau)^2},\\
a_3^2 e^{\phi} &= \frac{L_1 \sqrt{p_1}\, e^{(\frac{N-p_1}{2})\tau}}{\sinh(p_1\tau)^2},\\
a_4^2 e^{\phi} &= \frac{8(L_1B)^4\, e^{(N-2p_1)\tau}}{9\sinh(p_1\tau)^6},
\end{aligned} \tag{111}$$

where L_1 and p_1 are constants. From the initial value equation (19) we have the following restriction on p_1:

$$3N^2 - 29p_1^2 + 2Np_1 = 0. \tag{112}$$

4.25 Lie algebras $A_{4,9}^b$ $(b \neq \pm1, -2)$

We obtain the potentials V_i as follows:

$$\begin{aligned}
&V_1 = \frac{a_1^2}{2a_2^2a_3^2} - \frac{2(b+1)^2}{a_4^2},\ \ V_2 = -\frac{a_1^2}{2a_2^2a_3^2} - \frac{2(b+1)}{a_4^2},\\
&V_3 = -\frac{a_1^2}{2a_2^2a_3^2} - \frac{2b(b+1)}{a_4^2},\ \ V_4 = -\frac{2(b^2+b+1)}{a_4^2}.
\end{aligned} \tag{113}$$

To facilitate calculations we will assume for the moment that we also have $b \neq \pm1, -2$ and present the $A_{4,9}^{b=\pm1,2}$ models separately in the next subsection. In this case the constraint equations (10) impose the following restriction:

$$a_1^{b+1} a_2 a_3^b = B a_4^{2(b+1)}, \tag{114}$$

where B is constant. Equations (18) have the following forms:

$$\begin{aligned}
&\ln(a_1^2 e^{\phi})'' + \left(a_1^4 a_4^2 - 4(b+1)^2 a_1^2 a_2^2 a_3^2\right) e^{2\phi} = 0,\\
&\ln(a_2^2 e^{\phi})'' - \left(a_1^4 a_4^2 + 4(b+1) a_1^2 a_2^2 a_3^2\right) e^{2\phi} = 0,\\
&\ln(a_3^2 e^{\phi})'' - \left(a_1^4 a_4^2 + 4b(b+1) a_1^2 a_2^2 a_3^2\right) e^{2\phi} = 0,\\
&\ln(a_4^2 e^{\phi})'' - 4\left(b^2+b+1\right) a_1^2 a_2^2 a_3^2 e^{2\phi} = 0.
\end{aligned} \tag{115}$$

It follows that for $b \neq -2$ and $b \neq \pm 1$ the most general solution for these equations which satisfies the above constraint is:

$$\begin{aligned}
a_1^2 e^{\phi} &= \frac{L_1\, e^{p_1 \tau}}{\sinh(p_1\tau)^{\frac{4b^2+10b+4}{7b^2+13b+7}}}, \\
a_2^2 e^{\phi} &= \frac{L_2\, e^{\left(-\frac{b+2}{b-1}N+\frac{4b+5}{b-1}p_1\right)\tau}}{\sinh(p_1\tau)^{\frac{2b^2+8b+8}{7b^2+13b+7}}}, \\
a_3^2 e^{\phi} &= \frac{L_3\, e^{\left(-\frac{5b^2+14b+8}{(b+2)(b-1)}p_1+\frac{2b+1}{b-1}N\right)\tau}}{\sinh(p_1\tau)^{\frac{8b^2+8b+2}{7b^2+13b+7}}}, \\
a_4^2 e^{\phi} &= \frac{L_4\, e^{(N-2p_1)\tau}}{\sinh(p_1\tau)^{\frac{6(b^2+b+1)}{7b^2+13b+7}}},
\end{aligned} \tag{116}$$

where L_1 and p_1 are constants and the coefficient L_2, L_3 and L_4 are obtained as follows:

$$\begin{aligned}
L_2 &= \left(\frac{(3/2)^b L_1^{2b+3}}{B^2 p_1^{2(b+1)}} \frac{(7b^2+13b+7)^{b+2}}{(2(b^2+b+1))^{2(b+1)}}\right)^{\frac{1}{b-1}}, \\
L_3 &= \left(\frac{(2/3) B^2 p_1^{4b+2}}{L_1^{3b+2}} \frac{(2(b^2+b+1))^{2(b+1)}}{(7b^2+13b+7)^{2b+1}}\right)^{\frac{1}{b-1}}, \\
L_4 &= \frac{2(b^2+b+1)}{(7b^2+13b+7)} \frac{p_1^2}{L_1^2}.
\end{aligned} \tag{117}$$

From the initial value equation (19) we have the following restriction on p_1:

$$\begin{aligned}
&\left(\frac{3(7b^2+13b+7)}{(b-1)^2} - \frac{4(5b^2+8b+5)}{(7b^2+13b+7)}\right) p_1^2 + \left(\frac{3(b^2+b+1)}{(b-1)^2}\right) N^2 \\
&\quad - \frac{2(7b^2+13b+7)}{(b-1)^2} N p_1 = 0.
\end{aligned} \tag{118}$$

4.25.1 Lie algebras $A_{4,9}^b$ $(b = \pm 1, -2)$

i) For $b = 1$ the constraint equation (114) may be written as

$$a_1^2 a_2 a_3 = B a_4^4, \tag{119}$$

so that Eqs.(115) reduce to

$$\begin{aligned}
&\ln(a_1^2 e^{\phi})'' + \left(a_1^4 a_4^2 - 16 a_1^2 a_2^2 a_3^2\right) e^{2\phi} = 0, \\
&\ln(a_2^2 e^{\phi})'' - \left(a_1^4 a_4^2 + 8 a_1^2 a_2^2 a_3^2\right) e^{2\phi} = 0, \\
&\ln(a_3^2 e^{\phi})'' - \left(a_1^4 a_4^2 + 8 a_1^2 a_2^2 a_3^2\right) e^{2\phi} = 0, \\
&\ln(a_4^2 e^{\phi})'' - 12 a_1^2 a_2^2 a_3^2 e^{2\phi} = 0.
\end{aligned} \tag{120}$$

The general solution to these equations are

$$
\begin{aligned}
a_1^2e^{\phi} &= \frac{L_1\,e^{\frac{N}{3}\tau}}{\sinh^{\frac{2}{3}}(p_1\tau)},\\
a_2^2e^{\phi} &= \frac{L_2\,e^{p_2\tau}}{\sinh^{\frac{2}{3}}(p_1\tau)},\\
a_3^2e^{\phi} &= \frac{L_3\,e^{(\frac{2N}{3}-p_2)\tau}}{\sinh^{\frac{2}{3}}(p_1\tau)},\\
a_4^2e^{\phi} &= \frac{L_4\,e^{\frac{N}{3}\tau}}{\sinh^{\frac{2}{3}}(p_1\tau)},
\end{aligned}
\tag{121}
$$

where L_1, p_1, p_2 are constants and the coefficient L_2, L_3 and L_4 are obtained as follows:

$$
\begin{aligned}
L_1 &= (32B^2)^{\frac{1}{9}}(\frac{p_1}{3})^{\frac{2}{3}},\\
L_3 &= \frac{1}{36L_2}(\frac{16}{B^2})^{\frac{1}{9}}(3p_1^2)^{\frac{2}{3}},\\
L_4 &= \frac{(p_1/3)^{\frac{2}{3}}}{(2B^4)^{\frac{1}{9}}}.
\end{aligned}
\tag{122}
$$

From the initial value equation (19) we have the following restriction on p_2:

$$54p_2^2 - 58N^2 - 72Np_2 - 27N = 0. \tag{123}$$

ii) For $b = -1$ the constraint equation (114) may be written as

$$a_1a_2 = a_3, \tag{124}$$

and equations (115) get the following forms:

$$
\begin{aligned}
\ln(a_1^2e^{\phi})'' + a_1^4a_4^2e^{2\phi} &= 0,\\
\ln(a_2^2e^{\phi})'' - a_1^4a_4^2e^{2\phi} &= 0,\\
\ln(a_3^2e^{\phi})'' - a_1^4a_4^2e^{2\phi} &= 0,\\
\ln(a_4^2e^{\phi})'' - 4a_1^2a_2^2a_3^2e^{2\phi} &= 0,
\end{aligned}
\tag{125}
$$

which admit the following general solution:

$$
\begin{aligned}
a_1^2e^{\phi} &= L_1\,e^{p_1\tau}\sinh^2(p_1\tau),\\
a_2^2e^{\phi} &= \sqrt{\frac{3}{2L_1}}\frac{p_1\,e^{\left(\frac{N-p_1}{2}\right)\tau}}{\sinh^2(p_1\tau)},\\
a_3^2e^{\phi} &= \sqrt{\frac{3}{2L_1}}\frac{p_1\,e^{\left(\frac{N-p_1}{2}\right)\tau}}{\sinh^2(p_1\tau)},\\
a_4^2e^{\phi} &= \frac{2p_1^2\,e^{(N-2p_1)\tau}}{L_1^2\sinh^2(p_1\tau)},
\end{aligned}
\tag{126}
$$

where L_1 and p_1 are constants. From the initial value equation (19) we have the following restriction on p_1:

$$3N^2 - 29p_1^2 - 2Np_1 = 0. \tag{127}$$

iii) For case $b = -2$ the equations (114, 115) can be reduce to

$$a_2 a_4^2 = B a_1 a_3^2, \tag{128}$$

and

$$\begin{aligned}
&\ln(a_1^2 e^{\phi})'' + \left(a_1^4 a_4^2 - 4a_1^2 a_2^2 a_3^2\right) e^{2\phi} = 0,\\
&\ln(a_2^2 e^{\phi})'' - \left(a_1^4 a_4^2 - 4a_1^2 a_2^2 a_3^2\right) e^{2\phi} = 0,\\
&\ln(a_3^2 e^{\phi})'' - \left(a_1^4 a_4^2 + 8a_1^2 a_2^2 a_3^2\right) e^{2\phi} = 0,\\
&\ln(a_4^2 e^{\phi})'' - 12a_1^2 a_2^2 a_3^2 e^{2\phi} = 0,
\end{aligned} \tag{129}$$

which admit the following solutions:

$$\begin{aligned}
a_1^2 e^{\phi} &= L_1\, e^{p_1 \tau},\\
a_2^2 e^{\phi} &= (\frac{L_1 B^2}{16})^{\frac{1}{3}} e^{p_1 \tau},\\
a_3^2 e^{\phi} &= (\frac{2}{27A^4B^2})^{\frac{1}{3}} \frac{p_1^2\, e^{(N-2p_1)\tau}}{\sinh^2(p_1\tau)},\\
a_4^2 e^{\phi} &= \frac{2p_1^2\, e^{(N-2P_1)\tau}}{3L_1^2 \sinh^2(p_1\tau)},
\end{aligned} \tag{130}$$

where L_1 and p_1 are constants. From the initial value equation (19) we have the following restriction on p_1:

$$N^2 - p_1^2 + 2Np_1 = 0. \tag{131}$$

4.26 Lie algebras $A_{4,9}^0$

We obtain the potentials V_i as follows:

$$V_1 = \frac{a_1^2}{2a_2^2 a_3^2} - \frac{2}{a_4^2},\ \ V_2 = -\frac{a_1^2}{2a_2^2 a_3^2} - \frac{2}{a_4^2},\ \ V_3 = -\frac{a_1^2}{2a_2^2 a_3^2},\ \ V_4 = -\frac{2}{a_4^2}. \tag{132}$$

In this case the constraint equations (10) impose the following restriction:

$$a_1 a_2 = B a_4^2, \tag{133}$$

where B is constant. Equations (18) have the following forms:

$$\begin{aligned}
&\ln(a_1^2 e^{\phi})'' + \left(a_1^4 a_4^2 - 4a_1^2 a_2^2 a_3^2\right) e^{2\phi} = 0,\\
&\ln(a_2^2 e^{\phi})'' - \left(a_1^4 a_4^2 + 4a_1^2 a_2^2 a_3^2\right) e^{2\phi} = 0,\\
&\ln(a_3^2 e^{\phi})'' - a_1^4 a_4^2 e^{2\phi} = 0,\\
&\ln(a_4^2 e^{\phi})'' - 4a_1^2 a_2^2 a_3^2 e^{2\phi} = 0,
\end{aligned} \tag{134}$$

which admit the following general solutions:

$$\begin{aligned}
a_1^2 e^{\phi} &= \frac{L_1\, e^{p_1 \tau}}{\sinh^{\frac{4}{7}}(p_1 \tau)}, \\
a_2^2 e^{\phi} &= \frac{4 p_1^4 B^2\, e^{(2N - 5p_1)\tau}}{49 L_1^5 \sinh^{\frac{8}{7}}(p_1 \tau)}, \\
a_3^2 e^{\phi} &= \frac{21 L_1^4 e^{(4p_1 - N)\tau}}{8 B^2 p_1^2 \sinh^{\frac{2}{7}}(p_1 \tau)}, \\
a_4^2 e^{\phi} &= \frac{2 p_1^2\, e^{(N - 2p_1)\tau}}{7 L_1^2 \sinh^{\frac{6}{7}}(p_1 \tau)},
\end{aligned} \tag{135}$$

where L_1 and p_1 are constants. From the initial value equation (19) we have the following restriction on p_1:

$$21 N^2 + 127 p_1^2 - 98 N p_1 = 0. \tag{136}$$

4.27 Lie algebras $A_{4,10}$

We obtain the potentials V_i as follows:

$$\begin{aligned}
&V_1 = -\frac{a_1^2}{2a_2^2 a_3^2}, \;\; V_2 = \frac{a_1^2}{2a_2^2 a_3^2} - \frac{a_2^2}{2a_3^2 a_4^2} + \frac{a_3^2}{2a_2^2 a_4^2}, \\
&V_3 = \frac{a_1^2}{2a_2^2 a_3^2} + \frac{a_2^2}{2a_3^2 a_4^2} - \frac{a_3^2}{2a_2^2 a_4^2}, \;\; V_4 = \frac{a_2^2}{2a_3^2 a_4^2} + \frac{a_3^2}{2a_2^2 a_4^2} - \frac{1}{a_4^2}.
\end{aligned} \tag{137}$$

such that there are no constraint equation. Equations (18) have the following forms:

$$\begin{aligned}
&\ln(a_1^2 e^{\phi})'' - a_1^4 a_4^2 e^{2\phi} = 0, \\
&\ln(a_2^2 e^{\phi})'' + \left(a_1^2 a_3^4 - a_1^2 a_2^4 + a_4^2 a_1^4\right) e^{2\phi} = 0, \\
&\ln(a_3^2 e^{\phi})'' + \left(a_1^2 a_2^4 - a_1^2 a_3^4 + a_4^2 a_1^4\right) e^{2\phi} = 0, \\
&\ln(a_4^2 e^{\phi})'' + \left(a_1^2 a_3^4 - 2 a_1^2 a_2^2 a_3^2 + a_1^2 a_2^4\right) e^{2\phi} = 0,
\end{aligned} \tag{138}$$

which admit the following general solution:

$$\begin{aligned}
a_1^2 e^{\phi} &= \frac{p_1}{\sqrt{L_4}\, \sinh(p_1 \tau)}, \\
a_2^2 e^{\phi} &= a_3^2 e^{\phi} = L_2 e^{p_2 \tau} \sinh(p_1 \tau), \\
a_4^2 e^{\phi} &= L_4 e^{N \tau},
\end{aligned} \tag{139}$$

where L_2, L_4, p_1 and p_2 are constants. From the initial value equation (19) we have the following restriction on p_1 and p_2:

$$p_1^2 - p_2^2 + N^2 = 0. \tag{140}$$

4.28 Lie algebras $A^a_{4,11}$

We obtain the potentials V_i as follows:

$$\begin{aligned}
&V_1 = \frac{a_1^2}{2a_2^2a_3^2} - \frac{8b^2}{a_4^2},\ \ V_2 = -\frac{a_1^2}{2a_2^2a_3^2} + \frac{a_2^2}{2a_3^2a_4^2} - \frac{a_3^2}{2a_2^2a_4^2} - \frac{4b^2}{a_4^2},\\
&V_3 = -\frac{a_1^2}{2a_2^2a_3^2} - \frac{a_2^2}{2a_3^2a_4^2} + \frac{a_3^2}{2a_2^2a_4^2} - \frac{4b^2}{a_4^2},\ \ V_4 = -\frac{a_2^2}{2a_3^2a_4^2} - \frac{a_3^2}{2a_2^2a_4^2} - \frac{6b^2-1}{a_4^2}.
\end{aligned} \tag{141}$$

The constraint equations (10) impose in this case the restriction

$$a_1^2a_2a_3 = Ba_4^4, \tag{142}$$

where B is a constant. Equations (18) have the following forms:

$$\begin{aligned}
&\ln(a_1^2e^{\phi})'' + \left(a_1^4a_4^2 - 16b^2a_1^2a_2^2a_3^2\right)e^{2\phi} = 0,\\
&\ln(a_2^2e^{\phi})'' - \left(a_3^4 - a_2^4 + a_1^2a_4^2 + 8b^2a_2^2a_3^2\right)a_1^2e^{2\phi} = 0,\\
&\ln(a_3^2e^{\phi})'' - \left(a_2^4 - a_3^4 + a_1^2a_4^2 + 8b^2a_2^2a_3^2\right)a_1^2e^{2\phi} = 0,\\
&\ln(a_4^2e^{\phi})'' - \left(a_2^4 + a_3^4 + (12b^2-2)a_2^2a_3^2\right)a_1^2e^{2\phi} = 0.
\end{aligned} \tag{143}$$

These equations admit the following general solutions:

$$\begin{aligned}
a_1^2e^{\phi} &= \frac{(\frac{C}{b})^{\frac{2}{3}}\, e^{\frac{N\tau}{3}}}{2\sinh^{\frac{2}{3}}(p_1\tau)},\\
a_2^2e^{\phi} = a_3^2e^{\phi} &= \frac{(\frac{1}{\sqrt{C}b})^{\frac{2}{3}}\, e^{\frac{N\tau}{3}}}{3\sinh^{\frac{2}{3}}(p_1\tau)},\\
a_4^2e^{\phi} &= \frac{(b^2\sqrt{C})^{\frac{1}{3}}\, e^{\frac{N\tau}{3}}}{\sqrt{6B}\sinh^{\frac{2}{3}}(p_1\tau)},
\end{aligned} \tag{144}$$

where p_1 is a constant and $C = \frac{4}{3}(2Ba^{\frac{4}{3}})^{\frac{1}{3}}$.
From the initial value equation (19) we have the following restriction on p_1:

$$N = \pm 2p_1. \tag{145}$$

4.29 Lie algebras $A_{4,12}$

We obtain the potentials V_i as follows:

$$\begin{aligned}
&V_1 = \frac{a_1^2}{2a_2^2a_4^2} - \frac{a_2^2}{2a_1^2a_4^2} - \frac{2}{a_3^2},\ \ V_2 = -\frac{a_1^2}{2a_2^2a_4^2} + \frac{a_2^2}{2a_1^2a_4^2} - \frac{2}{a_3^2},\\
&V_3 = -\frac{2}{a_3^2},\ \ V_4 = -\frac{a_1^2}{2a_2^2a_4^2} - \frac{a_2^2}{2a_1^2a_4^2} + \frac{1}{a_4^2}.
\end{aligned} \tag{146}$$

In this case the constraint equations (10) impose the following restriction:

$$a_1 a_2 = B a_3^2, \tag{147}$$

where B is a constant. Equations (18) have the following forms:

$$\begin{aligned}
&\ln(a_1^2 e^{\phi})'' + \left(a_1^4 a_3^2 - a_2^4 a_3^2 - 4a_1^2 a_2^2 a_4^2\right) e^{2\phi} = 0,\\
&\ln(a_2^2 e^{\phi})'' - \left(a_1^4 a_3^2 - a_2^4 a_3^2 + 4a_1^2 a_2^2 a_4^2\right) e^{2\phi} = 0,\\
&\ln(a_3^2 e^{\phi})'' - 4a_1^2 a_2^2 a_4^2\, e^{2\phi} = 0,\\
&\ln(a_4^2 e^{\phi})'' - \left(a_1^4 a_3^2 + a_2^4 a_3^2 - 2a_1^2 a_2^2 a_3^2\right) e^{2\phi} = 0,
\end{aligned} \tag{148}$$

which admit the following general solution:

$$\begin{aligned}
a_1^2 e^{\phi} &= a_2^2 e^{\phi} = \frac{p_1\, e^{(\frac{N-p_4}{2})\tau}}{2\sqrt{L_4}\sinh(p_1\tau)},\\
a_3^2 e^{\phi} &= \frac{p_1\, e^{(\frac{N-p_4}{2})\tau}}{2B\sqrt{L_4}\sinh(p_1\tau)},\\
a_4^2 e^{\phi} &= L_4 e^{p_4\tau},
\end{aligned} \tag{149}$$

where L_4, p_1 and p_4 are constants. From the initial value equation (19) we have the following restriction on p_1 and p_4:

$$3N^2 - 12p_1^2 + 3p_4^2 - 2Np_4 = 0. \tag{150}$$

5 Example $VII_0 \oplus R$ as a 4+1 dimensional cosmology coupled to matter with negative pressure (accelerating universe)

In this section we apply one of the exact solutions of the previous section (e.g $VII_0 \oplus R$) as a physical cosmological model. In Ref. [8], it has been shown that the equations (2-4) are equivalent to the following Einstein equation

$$\tilde{R}_{\mu\nu} - \frac{1}{2}\tilde{R}\tilde{g_{\mu\nu}} = \kappa^2 (T^{(\phi)}_{\mu\nu} + T^{(H)}_{\mu\nu}) \tag{151}$$

with the metric tensor

$$\tilde{g_{\mu\nu}} = e^{\phi} g_{\mu\nu}, \tag{152}$$

and the following energy-momentum tensor for ϕ and H

$$\begin{aligned}
\kappa^2 T^{(\phi)}_{\mu\nu} &= \frac{1}{2}\left(\partial_\mu\phi\partial_\nu\phi - \frac{1}{2}(\partial\phi)^2\tilde{g_{\mu\nu}} - \Lambda e^{-\phi}\tilde{g_{\mu\nu}}\right),\\
\kappa^2 T^{(H)}_{\mu\nu} &= \frac{1}{4}\left(H_{\mu\kappa\lambda}H_{\nu}^{\kappa\lambda} - \frac{1}{6}H^2\tilde{g_{\mu\nu}}\right).
\end{aligned} \tag{153}$$

Now, we obtain the components of above tensors and Einstein equation for the example $VII_0 \oplus R$ of subsection 4. For this propose, we must obtain the metric $g_{\mu\nu}$ as functions of coordinate time t. From relations (13), (15) and (54) we obtain the following relation for τ as a function of t:

$$\tau = (\frac{1}{\gamma}) \ln(\frac{\gamma t}{\sqrt{L_1 L_2 L_3}}), \tag{154}$$

where $\gamma = \frac{(p_1+p_3+p_4-2N)}{2}$. After substituting (154) in relation (6) and using results of table 1 the metric tensor can be obtain as follows:

$$ds^2 = -dt^2 + a_1(t)^2 \left((dx^1)^2 + (dx^2)^2\right) + a_3(t)^2 (dx^3)^2 + a_4(t)^2 (dx^4)^2. \tag{155}$$

Using above equation one can calculate the non-zero components of energy-momentum tensor $T^{(\phi)}_{\mu\nu}$ for the $VII_0 \oplus R$ model as follows[4]:

$$\begin{aligned}
\kappa^2 T_{00} &= \frac{N^2}{2\gamma^2 t^2}\left(1 + \frac{1}{2}(\frac{\gamma t}{\sqrt{L_1 L_2 L_3}})^{\frac{N}{\gamma}}\right), \\
\kappa^2 T_{11} = \kappa^2 T_{22} &= -\sqrt{L_1}\frac{N^2}{4\gamma^2 t^2}\left(\frac{\gamma t}{\sqrt{L_1 L_2 L_3}}\right)^{\frac{p_1}{2\gamma}}, \\
\kappa^2 T_{33} &= -L_2 \frac{N^2}{4\gamma^2 t^2}\left(\frac{\gamma t}{\sqrt{L_1 L_2 L_3}}\right)^{\frac{p_3}{\gamma}}, \\
\kappa^2 T_{44} &= -L_3 \frac{N^2}{4\gamma^2 t^2}\left(\frac{\gamma t}{\sqrt{L_1 L_2 L_3}}\right)^{\frac{p_4}{\gamma}}.
\end{aligned} \tag{156}$$

Now by comparing the components of $T_{\mu\nu}$ with the following perfect fluid energy-momentum tensor:

$$T = \begin{pmatrix} -\rho(t) & 0 & 0 & 0 & 0 \\ 0 & P(t) & 0 & 0 & 0 \\ 0 & 0 & P(t) & 0 & 0 \\ 0 & 0 & 0 & P(t) & 0 \\ 0 & 0 & 0 & 0 & P(t) \end{pmatrix} \tag{157}$$

and assuming $\left(\kappa = 1, \frac{p_1}{2} = p_3 = p_4, \sqrt{L_1} = L_2 = L_3\right)$ and using (55), one can obtain the density and pressure quantities of the matter and scale factors as follows:

$$\begin{aligned}
\rho(t) &= -\frac{1}{2}\left(\frac{9}{t^2} + \frac{N^3}{6L_3^6}t\right), \\
P(t) &= -\frac{N^2}{4L_3^3}
\end{aligned} \tag{158}$$

$$a_i(t) = \sqrt{\frac{3L_3^3}{Nt}}, \qquad (i = 1, 2, 3, 4). \tag{159}$$

---

[4] Note that for our examples of section 4, $H_{\mu\nu\lambda} = 0$ and consequently $T^{(H)}_{\mu\nu} = 0$

We examine our model with a cosmological model of dark energy. Thus, from equation (158) we insist that the our model corresponds to the equation of state for dark energy as follows:

$$P(t) = -\rho(t), \tag{160}$$

So, by taking condition (160) and assuming $N = L_3^2$ we can get

$$H_0 = \frac{1}{t} = \frac{1}{81}\left(-54 + 6\sqrt{6L_3^3 + 81}\right)^{\frac{1}{3}} - \frac{L_3}{3\left(-54 + 6\sqrt{6L_3^3 + 81}\right)^{\frac{1}{3}}}, \tag{161}$$

where H_0 is the Hubble parameter at epoch that the universe accelerates in the context of cosmological constant regime. We can fix the parameter L_3 in our model so that leads to accelerating universe (see for example [20]). Also, we can determine another cosmological parameter so-called the deceleration parameters which has the following negative values

$$q_i(t) = -\frac{\ddot{a}_i(t)a_i(t)}{\dot{a}_i(t)^2} = -3. \tag{162}$$

From the above equation it can be seen we have $q < 0$. Thus, our model predicts an accelerating universe.

6 Conclusion

As a consequences let us examine the effect of Abelian target space duality on the above backgrounds. Consider the following sigma model action:

$$S = \int_\Sigma d^2\xi \left(g^{\alpha\beta}\partial_\alpha x^\mu \partial_\beta x^\nu G_{\mu\nu} + \varepsilon^{\alpha\beta} B_{\mu\nu}\partial_\alpha x^\mu \partial_\beta x^\nu + R^{(2)}\phi\right) \tag{163}$$

such that the equations (2-4) are the zeros of its one loop beta function anomaly. In the above equation g, $R^{(2)}$ are metric and scalar curvature of the two dimensional world sheet Σ respectively. In order to obtain the dual background it is enough to apply the Buscher's duality transformations [21] in the isometry direction.

The isometry directions of the $4+1$ dimensional homogenous anisotropic backgrounds are written in the table 1. Here we consider an example for $A_{4,8} \cong H_4$ Lie algebra. One can obtain all other dual backgrounds in the same way by using Buschers's duality [21]. For the Lie algebra $A_{4,8}$ the metric on the $4+1$ dimensional space-time is obtained as follows:

$$ds^2 = -dt^2 + a_1^2 dx_1^2 + 2a_1^2 x_3 dx_1 dx_2 + \left(a_1^2 x_3^2 + a_2^2 e^{2x_4}\right) dx_2^2 + a_3^2 e^{-2x_4} dx_3^2 + a_4^2 dx_4^2 \tag{164}$$

This metric has two isometry directions x_1 and x_2, such that we have $[X_1, X_2] = 0$. Now by use of the Buscher's duality transformation [21] in the x_1 direction one can obtain the following dual background:

$$\left\{\begin{array}{l} \tilde{G}_{11} = \frac{1}{a_1^2},\ \tilde{G}_{12} = \tilde{G}_{13} = \tilde{G}_{14} = 0, \\ \\ \tilde{G}_{22} = a_2^2 e^{2x_4},\ \tilde{G}_{23} = \tilde{G}_{24} = 0, \\ \\ \tilde{G}_{33} = a_3^2 e^{-2x_4},\ \tilde{G}_{34} = 0, \\ \\ \tilde{G}_{44} = a_4^2, \end{array}\right. \qquad \left\{\begin{array}{l} \tilde{B}_{12} = x_3,\ \tilde{B}_{13} = \tilde{B}_{14} = 0, \\ \\ \tilde{B}_{23} = \tilde{B}_{24} = 0,\quad \tilde{B}_{34} = 0, \end{array}\right. \tag{165}$$

$$H_{123} = 1,\ \tilde{\phi} = \phi(t) - \frac{1}{2}\ln a_1(t). \tag{166}$$

We see that in the dual background the antisymmetric tensor is nonzero. In this way, in order to obtain the solutions of equations of motions (2-4) for this backgrounds, it is enough to use of duality transformations. In the similar way one can obtain other dual backgrounds by considering isometry directions mentioned in the table 1 and using Buscher's duality transformations in these directions.

Acknowledgment

The authors is grateful to F. Darabi and K. Atazadeh for useful discussion and recommendation.

References

[1] R. Brandenberger and C. Vafa, Nucl. Phys. B, 316 (1989) 391.

[2] E. S. Fradkin and A. Tseytlin, Nucl. Phys. B. 261 (1985) 1; C. Callan, D. Friedan, E. Martinec and M. Perry; Nucl. Phys. B. 262 (1985) 593.

[3] M. Gasperini and G. Veneziano, Phys. Rep. 373 (2003) 1; see also J. E. Lidsey, D. wands and E. J. Copeland, Phys. Rep. 337 (2000) 343.

[4] V. Bozza, M. Gasperini, M. Giovannini and G. Veneziano, Phys. Lett. B, 543 (2002) 14.

[5] M. Gasperini, M. Giovannini and G. Veneziano, Nucl. Phys. B, 694 (2004) 206.

[6] E. J. Copeland, A. Lahiri and D. Wands. Phys. Rev. D, 50 (1994) 4868; Phys. Rev. D, 51 (1994) 1569.

[7] J. D. Barrow and K. E. Kunze, Phys. Rev. D, 55 (1997) 623.

[8] N. A. Batakis and A. A. Kehagias; Nucl. Phys B. 449 (1995) 248-264.

[9] M. Gasperini and R. Ricci, *Homogenous conformal string backgrounds*, Class. Quanmum. Grav. 12 (1995) 677-688.

[10] J. D. Barrow and K. E. Kunze, Phys. Rev. D,6 (1997) 741; D. Clancy, A. Feinstein, J. Lidsey and R. Tavakol, Pys. Rev. D, 63 (2001) 064002.

[11] G. F. R. Ellis and M. A. H. MacCallum, Commun. Math. Phys. 12 (1969) 108.

[12] S. Hervik, Quantum. Grav. 19 (2002) 5409-5427.

[13] L. D. Landau and E. M. Lifshits, *The Classical Theory of Fields*(Pergamon, 1987).

[14] T. Christodoulakis, S. Hervik, G. O. Papadopoulos, J. Phys. A: Math. Gen. 37 (2004) 4039-4058.

[15] G. M. Mubarakzyanov; Isv. Vyss. Uch. Zav. Mat, 4 (1963) 104-116.

[16] A. Z. Petrov, *Einstein spaces*, Pergamon Press (1961).

[17] J. Patera and P. Winternitz; J. Math .Phys, 18 (1977) 1449-1455.

[18] M. A. H. MacCallum, *On the classification of the real four-dimensional Lie algebras* in *On Einsteins Path: Essays in Honor of Engelbert Schucking*, ed. A. L. Harvey (Springer-Verlag, New York, 1999), pp. 299317.

[19] M. A. Jafarizadeh and A. Rezaei-Aghdam, Phys. Lett. B, 458 (1999) 477.

[20] F. Darabi, Class. Quant. Grav, 20 (2003) 3385-3402.

[21] T. H. Buscher, Phys. Lett. B, 201 (1988) 466-472.